



NAC RECOMMENDATIONS FOR THE USE OF SOLVENT-DETERGENT PLASMA IN CANADA



SOLVENT-DETERGENT PLASMA SUBCOMMITTEE

**Solvent Detergent Plasma
Subcommittee Members:**

Alan Tinmouth, MD, FRCPC, MSc
Douglas Morrison, MD
Jason Quinn, MD
Katerina Pavenski, MD
Kathryn Webert, MD
Ryan Lett, MD

NAC Chair:

Alan Tinmouth, MD, FRCPC, MSc

Provincial Ministry Representative:

Thomas Smith (ON)

NAC Coordinator:

Harleen Kahlon
Sebastian Ye

Publication Date:

March 10, 2023

Date of Last Revision:

July 20, 2023

Cite As:

Vancouver

Tinmouth A, Morrison D, Quinn J, Pavenski K, Webert K, Lett R. NAC Recommendations for the Use of Solvent-Detergent Plasma in Canada [Internet]. Ottawa: National Advisory Committee on Blood and Blood Products; March 10, 2023 [updated 2023 07 20; cited YYYY MM DD]. Available from: <https://nacblood.ca/en/resource/nac-recommendations-solvent-detergent-plasma-sd-plasma>



TABLE OF CONTENTS

SOLVENT-DETERGENT PLASMA SUBCOMMITTEE	2
TABLE OF CONTENTS.....	3
ACRONYMS	4
BACKGROUND.....	5
INDICATIONS.....	6
SPECIAL POPULATIONS	6
DOSING CONSIDERATIONS	8
RECOMMENDATIONS FOR PREPARING AND INFUSING OF SD PLASMA	9
REFERENCES	11



ACRONYMS

S/D Plasma

Solvent-Detergent Human Plasma

CADTH

Canadian Agency for Drugs and Technologies in Health

FP

Frozen Plasma

TTP

Thrombotic Thrombocytopenic Purpura



BACKGROUND

On March 27, 2023, the restrictions on the use of Octaplasma Solvent-Detergent Human Plasma (S/D Plasma) in Canada (outside the province of Quebec) will be removed. Prior to this date, S/D Plasma was restricted to specific patient groups as recommended by the Canadian Agency for Drugs and Technologies in Health¹. The restrictions on the use of S/D Plasma were based on clinical evidence showing no difference in effectiveness as compared to regular frozen plasma (FP), and a substantial increase in cost associated with S/D Plasma as compared to FP². In July 2022, the Canadian Blood Services announced a transition to pathogen-reduced platelet and plasma components as an additional layer of safety to the blood supply system in Canada. The first phase in the implementation of pathogen-reduced plasma is the transition from FP and S/D Plasma with a target of 80% of all transfused plasma to be S/D Plasma by September 2023. The second phase (planned for 2024-25) is to replace the remaining 20% of FP with pathogen-reduced frozen plasma using the same technology (Intercept Blood System) currently in use to produce pathogen-reduced platelets.

S/D Plasma has been used in Canada since 2011 on a restricted basis, and in many European countries (Netherlands, Finland, Sweden, Norway, United Kingdom) as the primary plasma product with no concerns for either clinical efficacy or safety. S/D Plasma is made from large pools of plasma that undergo pathogen inactivation using a solvent-detergent process and is then divided into 200 mL units. As a result of the manufacturing process, S/D Plasma units have relatively standard or uniform levels of coagulation factors. The levels for all clotting factors are similar to those in FP with the exception of lower protein S and anti-plasmin levels. The pooling of units for S/D Plasma results in decreased allergic reactions³⁻⁶ and possibly a reduction in TRALI reactions⁷.

The transition to S/D Plasma represents a significant change in hospital transfusion practice in Canada. While S/D Plasma and FP have the same clinical indications, there are some differences in the products themselves and the packaging of the product, which are important for health care practitioners and hospital transfusion medicine services. Further information from Canadian Blood Services on S/D Plasma and the transition to this product can be found in the publication, [Solvent detergent \(S/D\) treated plasma \(Octaplasma\)](#) and [FAQ on S/D treated plasma \(Octaplasma\)](#)⁸.

This document from the National Advisory Committee on Blood and Blood Products (NAC) provides additional clinical recommendations regarding the indications, dosing, and the use in special populations.



INDICATIONS

The main indication for plasma transfusion is the replacement of deficient coagulation factor(s) in patients with active bleeding, or prior to a surgery or invasive procedures. Specific guidelines and recommendations for the use of plasma transfusion have been previously published^{9,10}.

In general, most patients who would be treated with FP can receive S/D Plasma, these would include the following indications:

- Bleeding patients or patients undergoing invasive procedures who require replacement of multiple plasma coagulation factors,
- Patients with massive transfusion with clinically significant coagulation abnormalities,
- Patients on warfarin who are bleeding or need to undergo an invasive procedure before vitamin K could reverse the warfarin effect, and where prothrombin complex concentrate is not available or is contraindicated,
- Patients with selected coagulation factor or with rare specific plasma protein deficiencies for which a more appropriate alternative therapy is not available,
- Preparation of reconstituted whole blood, and
- Patients with thrombotic thrombocytopenic purpura (TTP).

There are a limited number of specific contraindications to S/D Plasma:

- Patients with IgA deficiency and documented anti-IgA antibodies. These patients will also potentially have allergic reactions to regular FP and should only receive FP from IgA deficient donors. IgA deficiency alone (no anti-IgA antibodies) is not a contraindication as most patients with this relatively common deficiency do not form antibodies and will not have an adverse reaction to blood components¹¹, and
- Patients with severe deficiency of protein S. S/D Plasma has significantly lower levels of protein S as compared to FP, which may result in an increased risk of blood clots. If these patients with severe deficiencies of protein S require a plasma transfusion, they should receive FP.

SPECIAL POPULATIONS

(a) Pediatrics:

Based on limited data, S/D Plasma and FP should be considered equally effective in pediatric patients and can be used interchangeably. Two studies which included 91 pediatric patients receiving S/D Plasma did not identify any safety concerns¹²⁻¹³. A secondary analysis of a large multinational study identified 62 pediatric critical care patients who received S/D plasma; there was no difference in the reduction in the INR compared to patients receiving FP and S/D plasma was associated with reduced mortality¹⁴. Adoption of S/D plasma for routine use in other jurisdictions has also not identified any concerns with pediatric patients⁵.



(b) Neonatology:

Based on limited data, there is no reason to expect differences in clinical efficacy with S/D Plasma as compared to FP. A retrospective study included 41 neonate patients and reported no clinical or safety concerns with S/D Plasma¹⁵. One non-published study reported no adverse events in 55 neonates receiving S/D plasma¹⁶. The reduction in adverse transfusion reactions associated with S/D Plasma may be advantageous.

(c) Pregnancy:

Based on limited clinical data, S/D Plasma and FP can be considered equally effective in pregnant patients and can be used interchangeably. A retrospective study including 37 obstetrical patients reported no clinical or safety concerns¹⁶. Adoption of S/D plasma for routine use including obstetric and gynecology patients also did not identify any clinical concerns in gynecology or obstetrical patients⁵.

(d) Geriatrics:

S/D plasma and FP Plasma should be considered equally effective in geriatric patients and can be used interchangeably with FP. While there are no studies of evaluating S/D plasma specifically in geriatric patients, there are no physiologic reasons that S/D plasma would be less effective or have increased adverse effects in geriatric patients. The reduction in adverse events associated with S/D Plasma may be considered advantageous in this patient group who would be at greater risk of adverse outcomes associated with a transfusion reaction.

(e) Liver transplantation / Liver disease:

S/D Plasma and FP can be considered equally effective in patients undergoing liver transplantation or coagulopathy associated with liver disease and can be used interchangeably. Three randomized controlled trial studies (n=115) evaluated S/D Plasma in patients with coagulopathy associated with liver disease or liver transplant patients¹⁷⁻¹⁹. There were no differences in clinical efficacy or adverse events.

(f) TTP:

Clinical studies have shown that S/D plasma is effective and safe when used as part of plasmapheresis treatments in patients with TTP²⁰⁻²². No clinical studies have compared the effectiveness of S/D plasma to FP or cryosupernatant. In the 2 clinical trials evaluating cryosupernatant in TTP patients, there was no evidence of improved outcomes as compared to FP²³⁻²⁵. The reduction in adverse events and lower risk of transfusion transmitted infections with S/D plasma may be considered advantageous in patients receiving large volume plasma transfusions as part of plasmapheresis treatment.



DOSING CONSIDERATIONS

An important factor in the dosing of S/D Plasma is the smaller unit volumes. FP units, which are made from individual blood donations, vary in size with an average volume of 289 mLs (± 16 mLs)²⁵ and have more variability in coagulation factor levels as each unit is from an individual donor. Overall, the mean concentration of coagulation factors in S/D plasma and FP are similar²⁵, but as a pooled blood product, the concentrations are more uniform in S/D Plasma.

The correct dose of plasma is determined by either weight-based dosing (hemodynamically stable patients) or ratio-based dosing (massively bleeding patients):

- (1) For the correction of coagulation factor levels in patients with acute bleeding, or prior to surgery or invasive procedures, the recommended dose for FP is 10-15 mLs/kg^{9,27}, which is the volume required to increase coagulation factors levels above the minimum hemostatic threshold of 30%²⁸. Given the similarity in coagulation factor levels, this dose is also appropriate for S/D Plasma. In pediatric patients weighing less than 40 kg, the plasma is usually ordered in mLs/kg. For larger pediatric patients and adult patients, plasma is usually ordered in units. As S/D Plasma is supplied in a smaller standard 200 mL bag compared to an average 289 mL per unit of FP, a larger number of units of S/D Plasma may be required to deliver a therapeutic dose of plasma. Most adult patients (60-90 kg) will require 4-6 units of S/D Plasma or 3-4 units of FP to achieve hemostatic levels of coagulation factors. The prescribed dose of plasma should be guided by the clinical situation and coagulation results.
- (2) The NAC recommends that the number of plasma units prescribed in Massive Hemorrhage Protocols (MHPs) be the same whether S/D Plasma or FP is used. For patients with massive hemorrhage requiring transfusion support, resuscitation with fixed ratio of FP to Red Blood Cells (RBCs) is commonly used in MHPs²⁹. This practice is based on observational studies that report improved outcomes in military patients with massive hemorrhage secondary to trauma. While a high transfusion ratio of FP:RBC has been recommended in trauma patients requiring massive transfusion, the ideal ratio is unknown³⁰. In clinical practice, there is variability in the volumes of the FP and RBC units transfused, and often the desired ratio of FP:RBC units is not achieved. As a result, there is significant inherent variability in the actual volumes of plasma and red cells transfused. FP:RBC ratios of 1:1 to 1:3 have been recommended as sufficient to provide hemostatic levels of coagulation factors in trauma patients³¹. Additionally, no benefit has been demonstrated for higher FP:RBC ratios in the resuscitation of patients with non-traumatic massive hemorrhage, who represent the majority of MHP activations in the civilian setting. Therefore, based on the limited evidence available, the NAC does not recommend increasing the initial ratio of plasma to RBC units in MHPs when using S/D plasma despite the lower volume of S/D plasma units. However, as per recent recommendations, patients requiring massive transfusion require frequent monitoring of coagulation tests



and should be transitioned to laboratory guided administration of blood components as soon as possible²⁹.

Recommendations for Preparing and Infusing of SD Plasma

(1) Administration rate:

A maximum infusion rate of 1 mL/kg/min is recommended in the product monograph for Octaplasma. In most clinical situations, the infusion rate for plasma will be much lower than the maximum recommended infusion rate. For a 70 kg patient, the maximum rate would allow for an infusion of 1 unit of Octaplasma in 3 minutes. Infusion rates above this rate would only be seen when a rapid infuser, which can infuse at rates up to 750-1000 mL/min, is used.

The concern related to higher infusion rates is due to the citrate anticoagulant in Octaplasma and the potential for citrate toxicity. However, the risk of citrate toxicity is not different for Octaplasma as compared to regular FP. The concentration of citrate in Octaplasma is 17 mM/L which is similar to the calculated estimate for FP (17-25 mM/L) [Unpublished data, Canadian Blood Services]. Therefore, Octaplasma and FP have a similar risk for citrate toxicity during high volume rapid infusion.

Rapid infusions of plasma only occur in the setting of a massive hemorrhage. In this setting a fixed ratio of red blood cells and frozen plasma is recommended to decrease the coagulopathy associated with trauma and possibly decrease mortality. The risk of citrate toxicity will be reduced by ongoing rapid blood loss, which includes the loss of citrate, and giving added calcium as part of a massive hemorrhage protocol. Based on current clinical practice recommendations for trauma, infusing plasma at rates above 1 mL/kg would be considered standard practice where the benefit of the rapid infusion outweighs the potential risk for citrate toxicity. NAC recommends that Octaplasma and FP can be used interchangeably in this setting.

(2) Temperature for thawing and infusion:

The product monograph recommends that during thawing Octaplasma should not be warmed above 37°C. This follows current practice for FP and the temperature requirement does not represent a change in laboratory practice.

The rationale provided for this maximum temperature of 37°C as the optimal temperature for the constituents of blood is not clear. As measured by in-vitro coagulation studies, there is evidence for increased coagulation activity of whole blood with hyperthermia. However, the mechanisms suggested for the increased coagulation activity is related to tissue factor, which is almost exclusively extravascular, and not due to proteins found in plasma or the cellular components of whole blood³¹.

With regards to blood warmers and rapid infusion devices, there are no specific recommendations for temperature for Octaplasma or FP. Rapid infusers currently used to infuse blood products including plasma may be set at temperatures up to 41°C. Given the short period of infusion for most rapid infusion devices, the temperature of the plasma is unlikely to



significantly exceed 37°C. The exception is the Belmont infusion device, which heats the plasma through magnetic induction. However, even if the temperature of the plasma increases above 37°C, any increase in temperature will be transient and the infused plasma will very quickly equilibrate with the recipient's body temperature, thereby negating any potential adverse effects of a short temperature excursion. Importantly, any changes in coagulation protein activity due to increased temperatures would be seen in both FP and Octaplasma. Thus, the potential for any adverse impact on the function of coagulation proteins from using a blood warmer that increases the temperature of plasma above 37°C would be negligible, and they would be identical for Octaplasma and FP.

(3) Reconstituted Whole Blood:

Reconstituted whole blood is made by mixing plasma and red blood cells and is used primarily for neonatal exchange transfusion and priming cardiac bypass circuits. Reconstitution of whole blood using plasma is part of clinical practice of medicine. While the product monograph states that Octaplasma can be mixed with red blood cells and platelets, Health Canada regulations are unclear on this issue as Octaplasma is considered a drug and not a blood component. The use of either Octaplasma and FP for reconstituted whole blood is scientifically and medically appropriate, and both products should be considered interchangeable for this use.



REFERENCES

1. Optimal Therapy Recommendation for the use of Solvent/Detergent-Treated Human Plasma. [Internet] Ottawa; Canadian Agency for Drugs and Technologies for Health, May 2011 [cited 2023 02 24]. Available from: http://www.cadth.ca/media/pdf/SDPlasma_rec-report_e.pdf
2. Octaplas Compared with Fresh Frozen Plasma to Reduce the Risk of Transmitting Lipid-Enveloped Viruses: An Economic Analysis and Budget Impact Analysis. [Internet] Ottawa; Canadian Agency for Drugs and Technologies for Health, May 2011 [cited 2023 02 24]. Available from: https://www.cadth.ca/sites/default/files/pdf/SDPlasma_econo-analysis_e.pdf
3. Krusius T AM, Tuimala J. Introduction of Octaplas® in clinical use decreased the rate of severe adverse reactions. *Vox Sanguinis* 2010;99s1: P-1018
4. McGonigle AM, Patel EU, et al. Solvent detergent treated pooled plasma and reduction of allergic transfusion reactions. *Transfusion* 2020;60: 54-61.
5. Saadah NH, Schipperus MR, Wiersum-Osselton JC, et al. Transition from fresh frozen plasma to solvent/detergent plasma in the Netherlands: comparing clinical use and transfusion reaction risks. *Haematologica* 2020;105: 1158-65.
6. Cushing MM, Pagano MB, Jacobson J, et al. Pathogen reduced plasma products: a clinical practice scientific review from the AABB. *Transfusion* 2019;59: 2974-88.
7. Klanderman RB, van Mourik N, Eggermont D, et al. Incidence of transfusion-related acute lung injury temporally associated with solvent/detergent plasma use in the ICU: A retrospective before and after implementation study. *Transfusion* 2022;62: 1752-62.
8. FAQ: Solvent detergent (S/D) treated plasma (Octaplasma). Ottawa, Canadian Blood Services, 2022 [cited 2023 0301] Available from: <https://professionaleducation.blood.ca/en/transfusion/publications/faq-solvent-detergent-sd-treated-plasma-octaplasma>
9. O'Shaughnessy DF, Atterbury C, Bolton Maggs P, et al. British Committee for Standards in Haematology, Blood Transfusion Task Force. Guidelines for the use of fresh-frozen plasma, cryoprecipitate and cryosupernatant. *Br J Haematol.* 2004;126:11-28
Davidson JC, Rahim S, Hanks SE, et al. Society of Interventional Radiology Consensus Guidelines for the Periprocedural Management of Thrombotic and Bleeding Risk in Patients Undergoing Percutaneous Image-Guided Interventions-Part I: Review of Anticoagulation Agents and Clinical Considerations: Endorsed by the Canadian Association for Interventional Radiology and the Cardiovascular and Interventional Radiological Society of Europe. *J Vasc Interv Radiol.* 2019 Aug;30(8):1155-1167.
10. Khandelwal A, Clarke G, Goldman M. Anaphylactic transfusion reactions and IgA deficiency [Internet]. Ottawa: Canadian Blood Services; 2021 [cited 2023 02 24]. Available from: professionaleducation.blood.ca/en/transfusion/publications/anaphylactic-transfusion-reactions-and-iga-deficiency
11. Spinella PC, Borasino S, Alten J. Solvent/Detergent-Treated Plasma in the Management of Pediatric Patients Who Require Replacement of Multiple Coagulation Factors: An Open-Label, Multicenter, Post-marketing Study. *Front Pediatr.* 2020;8:572.



12. Josephson CD, Goldstein S, Askenazi D et al. Safety and tolerability of solvent/detergent-treated plasma for pediatric patients requiring therapeutic plasma exchange: An open-label, multicenter, postmarketing study. *Transfusion*. 2022;62:396-405
13. Camazine MN, Karam O, Colvin R, et al. Outcomes Related to the Use of Frozen Plasma or Pooled Solvent/Detergent-Treated Plasma in Critically Ill Children*. *Pediatric Critical Care Medicine*. 2017;18(5):e215–23
14. Chekrizova V and Murphy WG. Solvent-detergent plasma: use in neonatal patients, in adult and paediatric patients with liver disease and in obstetric and gynaecological emergencies. *Trans. Med*. 2006;16:85-91.
15. Canadian Product Monograph, Octaplasma®, May 14 2020, <https://www.octapharma.ca/api/download/x/9775dbca5e/octaplasma-pm-en-20200514.pdf>
16. Freeman JW, Williamson LM, Llewelyn C, et al. A randomized trial of solvent/detergent and standard fresh frozen plasma in the treatment of the coagulopathy seen during orthotopic liver transplantation. *Vox Sanguinis*. 1998;74(S1):225-229.
17. Williamson LM, Llewelyn CA, Fisher NC, et al. A randomized trial of solvent/detergent-treated and standard fresh-frozen plasma in the coagulopathy of liver disease and liver transplantation: SD-treated plasma in coagulopathy of liver disease. *Transfusion*. 1999;39(11-12):1227-1234.
18. Bindi ML, Miccoli M, Marietta M, et al. Solvent detergent vs. fresh frozen plasma in cirrhotic patients undergoing liver transplant surgery: a prospective randomized control study. *Vox Sang*. 2013;105:137-143
19. Scully M, Longair I, Flynn M, et al. Cryosupernatant and solvent detergent fresh-frozen plasma (Octaplas) usage at a single centre in acute thrombotic thrombocytopenic purpura. *Vox Sanguinis*. 2007
20. Edel E, Al-Ali HK, Seeger S, et al. Efficacy and Safety Profile of Solvent/Detergent Plasma in the Treatment of Acute Thrombotic Thrombocytopenic Purpura: A Single-Center Experience. *Transfusion Medicine and Hemotherapy : Offizielles Organ der Deutschen Gesellschaft für Transfusionsmedizin und Immunhamatologie*. 2010 ;37(1):13-19.
21. Harrison CN, Lawrie AS, Iqbal A, Hunter A, Machin SJ. Plasma exchange with solvent/detergent treated plasma of resistant thrombotic thrombocytopenic purpura. *Br J Haematol*. 1996;94:756–758.
22. Rock G, Shumak KH, Sutton DM, et al. Cryosupernatant as replacement fluid for plasma exchange in thrombotic thrombocytopenic purpura. Members of the Canadian Apheresis Group. *Br J Haematol*. 1996;94(2):383-6
23. Rock G, Anderson D, Clark W, et al. Canadian Apheresis Group; Canadian Association of Apheresis Nurses. Does cryosupernatant plasma improve outcome in thrombotic thrombocytopenic purpura? No answer yet. *Br J Haematol*. 2005;129(1):79-86.
24. Circular of information for the use of human blood Components: Plasma Components. Ottawa: Canada, November 2022 [cited 2023 03 01]. Available from: https://www.blood.ca/sites/default/files/IM-00004_Revision_1.pdf.



25. Sheffield WP, Bhakta V, Talbot K, et al. Quality of frozen transfusable plasma prepared from whole blood donations in Canada: an update. *Transfus Apher Sci.* 2013 Dec;49(3):440-6
26. Gupta A, Bigham M. Blood components. In: Clarke G, Chargé S, editors. *Clinical Guide to Transfusion* [Internet]. Ottawa: Canadian Blood Services, 2021 [cited 2023 03 01]. Chapter 2. Available from: <https://professionaleducation.blood.ca>.
27. Callum JL, Dzik WH.. The use of blood components prior to invasive bedside procedures: a critical appraisal. In: Mintz PD (ed), *Transfusion therapy: clinical principles and practice*, 3rd edn. Bethesda: AABB Press, 2010: 1–35
28. Callum JL, Yeh CH, Petrosoniak A, et al. A regional massive hemorrhage protocol developed through a modified Delphi technique. *CMAJ Open* 2019. DOI:10.9778/cmajo.20190042
29. da Luz LT, Shah PS, Strauss R, et al.. Does the evidence support the importance of high transfusion ratios of plasma and platelets to red blood cells in improving outcomes in severely injured patients: a systematic review and meta-analyses. *Transfusion.* 2019;59:3337-3349.
30. Jason A. Gregory, Sonny S. et al. Optimizing Transfusion Ratios in Massive Transfusion Protocols: An Argument Against the 1:1:1 Dogma and Approach to Trauma Resuscitation, *Laboratory Medicine*, 2015;46:e46–e52
31. Levi M. Hemostasis and Thrombosis in Extreme Temperatures (Hypo- and Hyperthermia). *Semin Thromb Hemost.* 2018 Oct;44(7):651-655.